

Conclusions

Cesium (^{137}Cs and ^{133}Cs) and Alkali Metals

The concentrations of the three stable alkali elements K, Rb, and ^{133}Cs and the activity concentration of ^{137}Cs have been determined in various components of Swedish forest – bulk soil, rhizosphere, soil-root interface fraction, fungal mycelium, and fungal sporocarps (Vinichuk et al. 2010b). The soil-root interface fraction is distinctly enriched with K and Rb, compared with bulk soil. Potassium concentration increases in the order bulk soil < rhizosphere < fungal mycelium < soil-root interface < fungal sporocarps, whereas, Rb concentration increases in the order bulk soil < rhizosphere < soil-root interface < fungal mycelium < fungal sporocarps. Cesium is generally evenly distributed within bulk soil, rhizosphere, and soil-root interface fractions, indicating no ^{133}Cs enrichment in these forest compartments.

The uptake of K, Rb, and ^{133}Cs during the entire transfer process between soil and sporocarps occurs against a concentration gradient. For all three alkali metals, the levels of K, Rb, and ^{133}Cs are at least one order of magnitude higher in sporocarps than in fungal mycelium. Potassium uptake appears to be regulated by fungal nutritional demands for this element and fungi have a high preference for the uptake of Rb and K than for Cs. According to the efficiency of uptaking by fungi, the three elements can be ranked in the order $\text{Rb}^+ > \text{K}^+ > \text{Cs}^+$, with a relative ratio of 100:57:32.

Although the mechanism of Cs uptake by fungi may be similar to that of Rb, the uptake mechanism for K appears to be different. The variability in isotopic (atom) ratios of $^{137}\text{Cs}/\text{K}$, $^{137}\text{Cs}/\text{Rb}$, and $^{137}\text{Cs}/^{133}\text{Cs}$ in the fungal sporocarps

suggests they are not species specific. The relationships between the concentration ratio $^{137}\text{Cs}/^{133}\text{Cs}$ and K, Rb, and ^{133}Cs in fungal sporocarps have wide and inconsistent variation, and the concentration of K, Rb, and ^{133}Cs in sporocarps appears to be independent of the $^{137}\text{Cs}/^{133}\text{Cs}$ isotopic ratio.

The study of *S. variegatus* sporocarps (Vinichuk et al. 2011a) sampled within 1 km² forest area with high ^{137}Cs fallout from the Chernobyl accident confirms ^{133}Cs and ^{137}Cs uptake is not correlated with uptake of K; whereas, the uptake of Rb is closely related to the uptake of ^{133}Cs . Furthermore, the variability in ^{137}Cs and alkali metals (K, Rb, and ^{133}Cs) among genotypes in local populations of *S. variegatus* is high and the variation appears to be in the same range as found in species collected at different localities. The variations in concentrations of K, Rb, and ^{133}Cs and ^{137}Cs activity concentration in sporocarps of *S. variegatus* appear to be influenced more by local environmental factors than by genetic differences among fungal genotypes.

Cs (^{137}Cs and ^{133}Cs), K and Rb in *Sphagnum* Plants

The distribution of ^{137}Cs within *Sphagnum* plants (down to 20 cm depth) (Vinichuk et al. 2010a) is similar to stable K, Rb, and ^{133}Cs . The ^{137}Cs activity concentrations and concentrations of K, Rb, and ^{133}Cs are highest in the uppermost 0-10 cm segments of *Sphagnum* (in the capitula and the subapical segments) and gradually decrease in older parts of the plant. Such distribution can be interpreted as dependent on the living cells of capitula and living green segments in the upper part of *Sphagnum*. The ^{137}Cs appears to be taken up and relocated by *Sphagnum* plants in similar ways to the stable alkali metals, as concentrations of K, Rb, and ^{137}Cs activity concentrations in *Sphagnum* segments are similar to the depth of about 16 cm, and display a slightly different pattern in the lower part of the plant.

Alkali Earth Metals Ca and Sr

Concentrations of Ca and Sr in soil fractions: bulk soil, rhizosphere, soil-root interface, fungal mycelium and fruit bodies of fungi are relatively high. The concentration of Ca and Sr is found to be higher in the rhizosphere fraction. Fungal sporocarps contained noticeably less calcium and strontium than mycelium and the mechanism of Ca and Sr exclusion appears efficient.

Fungal sporocarps varied in their ability to accumulate alkaline earth metals, although the concentration of calcium within fruit bodies of different species was several orders of magnitude higher than the concentration of strontium.

Calcium uptake correlates fairly well with Sr uptake in fungi: the Pearson correlation of Sr and Ca in fruit bodies of fungi is 0.904 (P-value = 0.000).

Transition Metals Cr, Co, Ni, Cu, Zn, Cd, Hg, and Pb

The capability of mycorrhizal fungi sporocarps to accumulate copper is higher than in mycelium: fungal sporocarps have statistically higher concentrations of Cu than the bulk soil (Vinichuk, 2012b, 2013). Zinc concentration in fungal mycelium is 2-fold higher than in bulk soil and 3-fold higher in sporocarps. The accumulation of cadmium is pronounced in both mycelium and fungal sporocarps: Cd concentration in mycelium is about 5 times higher than in bulk soil and about 2 times higher in sporocarps than in mycelium extracted from soil in the same plots where fungal sporocarps were sampled. Accumulation of Cd by fungal sporocarps is apparently species dependent.

Zinc concentration in sporocarps, and to a lesser extent in mycelium, depends on the concentration of Zn in the soil, whereas, the uptake of Cu and Cd by both sporocarps and mycelium does not correlate with their concentration in soil.

Thus, the uptake of Zn and Cu by fungi is balanced, implying similarities in the uptake mechanism.

The uptake of Cu, Zn, and Cd during the entire transfer process between soil and sporocarps occurs against a concentration gradient. For all three metals, the levels of Cu, Zn, and Cd in sporocarps is about two times higher than in fungal mycelium. Thus, fungi (mycelium and sporocarps) preferentially bioaccumulate Cd over Zn and Cu, and the bioconcentration values can be ranked in the order: $Cd^{2+} > Zn^{2+} > Cu^{2+}$, with a relative ratio of 100:41:38.

The concentration of Co and Ni in mycelium (Vinichuk, 2012a) is similar to the concentrations in other soil fractions (bulk soil, rhizosphere, soil-root interface); whereas, the concentrations in sporocarps are about 5-9 times lower than in mycelium. The concentration of Pb in mycelium is about 1.5 times lower than in other soil fractions (bulk soil, rhizosphere, soil-root interface) and about 50 times lower than in sporocarps. The mycorrhizal fungi (mycelium and sporocarps) only absorb Co, Ni, and Pb but do not accumulate these metals in tissue.

Fungal mycelium of upper forest soil layer may comprise 2.9-5.8% of chromium and 2.7-5.4% of nickel from their total content in soil. Fungal mycelium can also comprise between 4.6-9.1% of the total Cu in soil, 5.4-10.9% of the total Zn in the soil, and 15.5-31.7% of the total Cd in the soil. In fungal mycelium, the accumulation is estimated to be 3.5-6.9% for cobalt, 2.0-3.9% for Pb and 4.0-8.0% for Hg. In the mycelium of fungi in the upper (0-5 cm) layer of forest soil can be allocated 2.9-5.8% of the total amount of chromium in soil.

Semimetals (As)

No difference in As concentration in bulk soil and rhizosphere has been found, when soil-root interface fraction contains lower amounts of the elements (Unpublished data). The concentration of arsenic in mycelium was about the same as concentration in the soil and about 1.5 times higher in fruit bodies than in mycelium. Thus, arsenic does not appear to be accumulated by mycelium and only moderately by fruit bodies of fungi. Fungal mycelium accumulates between 3.0 and 6.1% of the total As in soil.

Actinides Th and U

Thorium concentration varies from of 0.0041 mg kg⁻¹ dry weight in fruit bodies of fungi to 1.45 mg kg⁻¹ dry weight fraction in the rhizosphere (Vinichuk, 2012c). Uranium concentration varies from 0.026 mg kg⁻¹ dry weight in fruiting bodies of fungi to 9.36 mg kg⁻¹ dry weight fraction in the rhizosphere. Both natural isotopes do not accumulate in either mushroom mycelium or their fruit bodies: bioconcentration ratios in fruit bodies are on average 0.006 for Th and 0.035 for U. Concentrations of Th and U isotopes in fruit bodies of fungi are about 270 times lower than in the bulk soil. The content of Th and U in mushroom mycelium and fruiting bodies depends on the concentration of these elements in the soil: as the concentrations of these elements in the soil increases, their content in mushrooms also increases. Mycelia of upper (0-5 cm) layers of the forest soil can comprise up to 5.0% of total thorium and 3.0% of total uranium content in soil.

References

- [1] Alonso, J.; Garcia, A. M.; Pérez-López, M.; Melgar, M. J. (2003). The concentrations and bioconcentration factors of copper and zinc in edible mushrooms. *Arch. Environ. Contam. Toxicol.* Vol. 44(2), 180-188. ISSN 0090-4341.
- [2] Adema, E.; Baaijens, G.; van Belle, J.; Rappoldt, A.; Grootjans, A. & Smolders, A. (2006). Field evidence for buoyancy-driven water flow in a *Sphagnum* dominated peat bog. *Journal of Hydrology*, Vol. 327, pp. 226-234, ISSN 0022-1694.
- [3] Baeza, A.; Hernández, S.; Guillén, F.; Moreno, J.; Manjón, J. L. & Pascual, R. (2004). Radiocaesium and natural gamma emitters in mushrooms collected in Spain. *Science of the Total Environment*, Vol. 318, No. 1-3, pp. 59-71, ISSN 0048-9697.
- [4] Baeza, A.; Guillén, J.; Hernández, S.; Salas, A.; Bernedo, M.; Manjón, J. & Moreno, G. (2005). Influence of the nutritional mechanism of fungi (mycorrhize/saprophyte) on the uptake of radionuclides by mycelium. *Radiochimica Acta*, Vol. 93, No. 4, pp. 233-238, ISSN 0033-8230.
- [5] Baeza, A. & Guillén, J. (2006). Influence of the soil bioavailability of radionuclides on the transfer of uranium and thorium to mushrooms. *Applied Radiation & Isotopes*, Vol. 64, No.9, pp. 1020-1026, ISSN 0969-8043.
- [6] Ban-nai T.; Yoshida S.; Muramatsu Y. & Suzuki A. (2005). Uptake of Radiocesium by Hypha of Basidiomycetes – Radiotracer Experiments. *Journal of Nuclear and Radiochemical Sciences*, Vol. 6, No.1, pp. 111-113, ISSN 1345-4749.
- [7] Basta, N. T.; Ryan, J. A.; Chaney, R. L. (2005). Trace element chemistry in residual-treated soil: key concepts and metal bioavailability. *Journal of Environmental Quality*, Vol. 34, No. 1, pp. 49-63, ISSN 0047-2425.
- [8] Bates, J. (1997). Effects of intermittent desiccation on nutrient economy and growth of two ecologically contrasted mosses. *Annals of Botany*, Vol.79, pp. 299-309, ISSN 0305-7364.

- [9] Berthelsen, B. O.; Olsen, R. A.; Steinnes, E. (1995). Ectomycorrhizal heavy metal accumulation as a contributing factor to heavy metal levels in organic surface soils. *Science of the Total Environment*, Vol. 170, No.1-2, pp. 141-149, ISSN 0048-9697.
- [10] Berthelsen, B. O.; Lamble, G. M.; MacDowell, A. A.; Nicholson, D. G. (2000). Analysis of Metal Speciation and Distribution in Symbiotic Fungi (Ectomycorrhiza) Studied by Micro X-ray Absorption Spectroscopy and X-ray Fluorescence Microscopy. In: G. Gorban, W. Venzel, & E. Lombi (Eds.), *Trace Elements in the Rhizosphere*, pp. 149-165, ISBN 0-8493-1535-2, Vienna, Austria.
- [11] Blanuša, M.; Antun, K. A.; Varnai, V-M.; Matek, S. M. (2001). Uptake of Cadmium, Copper, Iron, Manganese, and Zinc in Mushrooms (*Boletaceae*) from Croatian Forest Soil. *Journal of AOAC International*, Vol. 84, No.6, pp. 1964-1971, ISSN 1060-3271.
- [12] Borovička, J.; Kubrová J.; Rohovec, J.; Randa, Z.; Dunn, C. (2011). Uranium, thorium and rare earth elements in macrofungi: what are the genuine concentrations? *Biometals*, Vol. 24, No.5, pp. 837-845, ISSN 0966-0844.
- [13] Brzostowski, A.; Jarzyńska, G.; Kojta, A. K.; Wydmańska, D.; Falandysz, J. (2011). Variations in metal levels accumulated in Poison Pax (*Paxillus involutus*) mushroom collected at one site over four years. *Journal of Environmental Sciences*, Vol. 46, No.6, pp. 581-588. ISSN 1093-4529.
- [14] Brown, G. & Cummings, S. (2001). Potassium uptake and retention by *Oceanomonas baumannii* at low water activity in the presence of phenol. *FEMS Microbiology Letters*, Vol. 205, No. 1, pp. 37-41, ISSN 0378-1097.
- [15] Brun, C. B., Peltola, P., Åström, M. E., Johansson, M. B. (2010). Spatial distribution of major, trace and ultra trace elements in three Norway spruce (*Picea abies*) stands in boreal forests, Forsmark, Sweden. *Geoderma*, Vol. 159, pp. 252-261. ISSN 016-7061.
- [16] Brunner, I.; Frey, B. & Riesen, T. (1996). Influence of ectomycorrhization and cesium/potassium ratio on uptake and localization of cesium in Norway spruce seedlings. *Tree Physiology*, Vol. 16, pp. 705-711, ISSN 0829-318X.

- [17] Bunzl, K. & Kracke, W. (1989). Seasonal variation of soil-to-plant transfer of K and fallout $^{134,137}\text{Cs}$ in peatland vegetation. *Health Physics*, Vol.57, pp. 593-600, ISSN 0017-9078.
- [18] Burgess, T. I.; Maljczuk, N.; Grove, T. S. (1993). The ability of 16 ectomycorrhizal fungi to increase growth and phosphorus uptake of *Eucalyptus globulus* Labill. and *E. diversicolor* F. Muell. *Plant and Soil*, Vol. 153, No.2, pp. 155-164, ISSN 0032-079X.
- [19] Byrne, A. R.; Ravnik, V.; Kosta, L. (1976). Trace element concentrations in higher fungi. *Science of the Total Environment*, Vol. 6, No.1, pp. 65-78, ISSN 0048-9697.
- [20] Bystrzejewska-Piotrowska, G. & Bazala, M. (2008). A study of mechanisms responsible for incorporation of cesium and radiocaesium into fruitbodies of king oyster mushroom (*Pleurotus eryngii*). *Journal of Environmental Radioactivity*, Vol. 99, pp. 1185-1191, ISSN 0265-931X.
- [21] Bååth, E. (1980). Soil fungal biomass after clear-cutting of a pine forest in central Sweden. *Soil Biology & Biochemistry*, Vol. 12, No.5, pp. 495-500, ISSN 0038-0717.
- [22] Campbell, L.; Fisk, A.; Wang, X.; Köck, G.; D. Muir. (2005). Evidence for biomagnification of rubidium in freshwater and marine food webs. *Canadian Journal of Fisheries and Aquatic Science*, Vol. 62, pp. 1161-1167, ISSN 0706-652X.
- [23] Campos, J.; Tejera, N.; Sánchez, C. (2009). Substrate role in the accumulation of heavy metals in sporocarps of wild fungi, *Biometals*, Vol. 22, No.5, pp. 835-841, ISSN 0966-0844.
- [24] Campos, J.; De Toro, J.; Pérez de los Reyes, C.; Amorós, J.; García-Moreno, R. (2012). Lifestyle Influence on the Content of Copper, Zinc and Rubidium in Wild Mushrooms. *Applied and Environmental Soil Science*, Vol. 2012, Article ID 687160, 6 pages. ISSN 1687-7667.
- [25] Chao, J.; Chiu, C.; Lee, H. (2008). Distribution and uptake of ^{137}Cs in relation to alkali metals in a perhumid montane forest ecosystem. *Applied Radiation and Isotopes*, Vol.66, pp. 1287-1294, ISSN 0969-8043.
- [26] Collin-Hansen, C.; Yttri, K. E.; Andersen, R. A.; Berthelsen, B. O.; Steinnes, E. (2002). Mushrooms from two metal-contaminated areas in Norway: occurrence

- of metals and metallothionein-like proteins. *Geochemistry: Exploration, Environment, Analysis*, Vol. 2, No.2, pp. 121-130, ISSN 1467-7873.
- [27] Conway, E. & Duggan, F. (1958). A cation carrier in the yeast cell wall. *Biochemistry Journal*, Vol. 69, pp. 265-274, ISSN 0006-2960.
- [28] Clymo, R. (1963). Ion exchange in Sphagnum and its relation to bog ecology. *Annals of Botany*, (Lond.) Vol.27, pp. 309-324, ISSN 0305-7364.
- [29] Clymo, R. (1983). Peat. In: Gore, A. J. P. (Ed.), *Ecosystems of the world*. 4A. Mires: swamp, bog, fen and moor. General studies, 159-224, ISBN 0-444-42003-7, Elsevier, Amsterdam.
- [30] Clymo, R.; & Mackay, D. (1987). Upwash and downwash of pollen and spores in the unsaturated surface layer of *Sphagnum*-dominated peat. *New Phytologist*, Vol. 105, pp. 175-183, ISSN 0028-646X.
- [31] Dahlberg, A.; Nikolova, I.; Johanson, K. (1997). Intraspecific variation in ¹³⁷Cs activity concentration in sporocarps of *Suillus variegatus* in seven Swedish populations. *Mycological Research*, Vol. 101, pp. 545-551, ISSN 0953-7562.
- [32] Demirbaş, A. (2002). Metal ion uptake by mushrooms from natural and artificially enriched soils. *Food Chemistry*. Vol. 78(1), pp. 89-93, ISSN 0021-8561.
- [33] Dragović, S.; Nedić, O.; Stanković, S.; Bačić, G. (2004). Radiocaesium accumulation in mosses from highlands of Serbia and Montenegro: chemical and physiological aspects. *Journal of Environmental Radioactivity*, Vol.77, pp. 381-388, ISSN 0265-931X.
- [34] Drobner, U. & Tyler G. (1998). Conditions controlling relative uptake of potassium and rubidium by plants from soils. *Plant and Soil*, Vol.201, pp. 285-293, ISSN 0032-079X.
- [35] Elekes, C. C.; Busuioc, G.; Ionita, G. (2010). The Bioaccumulation of Some Heavy Metals in the Fruiting Body of Wild Growing Mushrooms. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, Vol. 38, No. 2, pp. 147-151, ISSN 0255-965X.
- [36] Enghag, P. (2000). *Jordens grund ännen och deras uppt äkt*. ISBN 9789175485904, Industrilitteratur. Stockholm.

- [37] Falandysz, J.; Kunito, T.; Kubota, R.; Gucia, M.; Mazur, A.; Falandysz, J. J.; Tanabe, S. (2008). Some mineral constituents of Parasol Mushroom (*Macrolepiota procera*). *Journal of Environmental Science and Health, Part B*, Vol. 43, pp. 187-192, ISSN 0360-1234.
- [38] Falandysz, J.; Frankowska, A.; Jarzyńska, G.; Dryżałowska, A.; Kojta, A. K.; Zhang, D. (2011). Survey on composition and bioconcentration potential of 12 metallic elements in King Bolete (*Boletus Edulis*) mushroom that emerged at 11 spatially distant sites. *Journal of Environmental Science and Health, Part B*, Vol.46, pp. 231-246, ISSN 0360-1234.
- [39] Falandysz, J.; Kojta, A.; Jarzyńska, G.; Drewnowska, M.; Dryżałowska, A.; Wydmańska, D.; Kowalewska, I.; Wacko, A.; Szlosowska, M.; Kannan, K.; Szefer, P. (2012). Mercury in bay bolete (*Xerocomus badius*): bioconcentration by fungus and assessment of element intake by humans eating fruiting bodies. *Food additives & contaminants. Part A, Chemistry, analysis, control, exposure & risk assessment*, Vol. 29, No.6, pp. 951-61, ISSN 1944-0049.
- [40] Fomina, M.; Hillier, S.; Charnock, J. M.; Melville, K.; Alexander, I. J.; Gadd, G. M. (2005). Role of oxalic acid overexcretion in transformations of toxic metal minerals by *Beauveria caledonica*. *Applied and Environmental Microbiology*, Vol.71, No.1, 371-381, ISSN 0099-2240.
- [41] Figueiredo, E.; Soares, M.; Baptista, P.; Castro, M.; Bastos, M. (2007). Validation of an Electrothermal Atomization Atomic Absorption Spectrometry Method for Quantification of Total Chromium and Chromium (VI) in Wild Mushrooms and Underlying Soils. *Journal of Agricultural and Food Chemistry*, Vol. 43, No.17, pp. 7192-7198, ISSN 0021-8561.
- [42] Galun, M.; Keller, P.; Malki, D.; Fedelstein, H.; Galun, E.; Siegel, S.; Siegel, B. (1983). Recovery of uranium (VI) from solution using precultured *Penicillium* biomass. *Water, Air, & Soil Pollution*, Vol. 20, pp. 221-232, ISSN 0049-6979.
- [43] Gaso, M.; Segovia, N.; Morton, O.; Cervantes, M.; Godinez, L.; Pe ña, P.; Acosta, E. (2000). ¹³⁷Cs and relationships with major and trace elements in edible mushrooms from Mexico. *Science of the Total Environment*, Vol. 262, No. 1-2, pp. 73-89, ISSN 0048-9697.

- [44] Gillett, A. & Crout, N. (2000). A review of ^{137}Cs transfer to fungi and consequences for modeling environmental transfer. *Journal of Environmental Radioactivity*, Vol. 48, pp. 95-121, ISSN 0265-931X.
- [45] Gorban, G. & Clegg, S. (1996). A conceptual model for nutrient availability in the mineral soil-root system. *Canadian Journal of Soil Science*, Vol.76, pp. 125-131, ISSN 1918-1841.
- [46] Greger M., (2004). Uptake of nuclides by plants. Report to SKB TR-04-14. 70 pp.
- [47] Gstoettner, E. & Fisher, N. (1997). Accumulation of cadmium, chromium, and zinc by the moss *Sphagnum papillosum* Lindle. *Water, Air, Soil Pollution*, Vol. 93, pp. 321-330, ISSN 0049-6979.
- [48] Gyuricza, V.; Dupré de Boulois, H.; Declerck, S. (2010). Effect of potassium and phosphorus on the transport of radiocaesium by arbuscular mycorrhizal fungi. *Journal of Environmental Radioactivity*, Vol.101, pp. 482-487, ISSN 0265-931X.
- [49] Hájek, T. (2008). Ecophysiological adaptations of coexisting *Sphagnum* mosses. PhD. thesis. University of South Bohemia, Faculty of Science, Czech Republic, 98 pp.
- [50] Högström, S. (1997). Habitats and increase of *Sphagnum* in the Baltic Sea island Gotland, Sweden. *Lindbergia*, Vol. 22, pp. 69-74, ISSN 0105-0761.
- [51] Holan, Z.; Volesky, B. (1995). Accumulation of cadmium, lead, and nickel by fungal and wood biosorbents, *Applied Biochemistry and Biotechnology*, Vol. 53, No.2, pp. 133-146, ISSN 0273-2289.
- [52] Horton, T. R. & Bruns, T. D. (2001). The molecular revolution in ectomycorrhizal ecology: peeking into the black-box. *Molecular Ecology*, Vol. 10, No.8, pp. 1855-1871, ISSN 1365-294X.
- [53] Horyna, J. & Řaná, Z. (1988). Uptake of radiocaesium and alkali metals by mushrooms. *Journal of Radioanalytical and Nuclear Chemistry*, Vol.127, pp. 107-120, ISSN 0236-5731.
- [54] Ismail, S. (1994). Distribution of Na, K, Rb, Cs and ^{137}Cs in some Austrian higher fungi. *Biology of Trace Element Research*, Vol.43-45, No.1, pp. 707-714, ISSN 01634984.

- [55] Johanson, K. J. & Bergström, R. (1994). Radiocaesium transfer to man from moose and roe deer in Sweden. *Science of the Total Environment*, Vol.157, pp. 309-316, ISSN 0048-9697.
- [56] Jonnalagadd, S.; Pienaar, D.; Haripersad, K. (2006). Elemental distribution in selected Agaricus and Rhizina mushrooms in South Africa. *Journal of Environmental Science and Health, Part A. Toxic/Hazardous Substances and Environmental Engineering*. Vol. 41, No 3. pp. 507-514, ISSN 1093-4529.
- [57] Kalač, P.; Burda, J.; Stašková, I. (1991). Concentrations of lead, cadmium, mercury and copper in mushrooms in the vicinity of a lead smelter. *Science of the Total Environment*, Vol. 105, pp. 109-119, ISSN 0048-9697.
- [58] Kalač P. (2001). A review of edible mushroom radioactivity. *Food Chemistry*, Vol.75, pp. 29-35, ISSN 0308-8146.
- [59] Kammerer, L.; Hiersche, L.; Wirth, E. (1994). Uptake of radiocaesium by different species of mushrooms. *Journal of Environmental Radioactivity*, Vol.23, pp. 135-150, ISSN 0265-931X.
- [60] Karadeniz, Ö. & Yaprak, G. (2007). Dynamic equilibrium of radiocesium with stable cesium within the soil-mushroom system in Turkish pine forest. *Environmental Pollution*, Vol. 148, No.1, pp. 316-324, ISSN 0269-7491.
- [61] Kirpichtchikova, T. A.; Manceau, A.; Spadini, L.; Panfili, F.; Marcus, M. A.; Jacquet, T. (2006). Speciation and solubility of heavy metals in contaminated soil using X-ray microfluorescence, EXAFS spectroscopy, chemical extraction, and thermodynamic modeling. *Geochimica et Cosmochimica Acta*, Vol. 70, No.9, pp. 2163-2190, ISSN 0016-7037.
- [62] Krznaric, E.; Wevers, J.; Colpaert, J. (2008). Heavy metal tolerant fungi play a key role in the protection of pines against combined Zn-Cd toxicity. 21st New Phytologist Symposium “The ecology of ectomycorrhizal fungi”, 10-12 December, 2008, Montpellier, France. Available from:
<http://www.newphytologist.org/mycorrhizal/21stNPS.pdf>.
- [63] Leake, J. & Read, D. (1997). Mycorrhizal fungi in terrestrial habitats. Chapter 18. In: *The Mycota, Vol. IV: Environmental and Microbial Relationship*, D. T. Wicklow & B. Söderström, (Ed.), 281-301, ISBN 978-3540-71839-0, Springer Verlag, Berlin.

- [64] Lepp, N. W.; Harrison, S. C. S.; Morrell, B. G. (1987). A role for *Amanita muscaria* L. in the circulation of cadmium and vanadium in a non-polluted woodland. *Environmental Geochemistry and Health*, Vol. 9, No. 3-4, pp. 61-64, ISSN 0269-4042.
- [65] Lundin, L.; Lode, E.; Stendahl, J.; Merklund, P.; Björkvald, L.; Thorstensson. (2004). Soils and site types in the Forsmark area, SKB-rapport, 102 p.
- [66] Marschner, H. (1995). *Mineral nutrition of higher plants*. Academic press, 2nded, ISBN 0-12-473543-6, London, UK.
- [67] Mascanzoni, D. (2009). Long-term transfer of ¹³⁷Cs from soil to mushrooms in a semi-natural environment. *Journal of Radioanalytical and Nuclear Chemistry*, Vol.282, pp. 427-431, ISSN 0236-5731.
- [68] Mattsson, S. & Lidén, K. (1975). ¹³⁷Cs in carpets of the forest moss *Pleurozium schreberi*, 1961-1973. *Oikos*, Vol.26, pp. 323-327, ISSN 0030-1299.
- [69] McGee, E.; Synnott, H.; Johanson, K.; Fawaris, B.; Nielsen, S. Horrill, A. D. (2000). Chernobyl fallout in a Swedish spruce forest ecosystem. *Journal of Environmental Radioactivity*, Vol.48, pp. 59-78, ISSN 0265-931X.
- [70] Meijer, R.; Aldenkamp, F.; Jansen, A. (1988). Resorption of cesium radionuclides by various fungi. *Oecologia*, Vol. 77, pp. 268-272, ISSN 0029-8549.
- [71] Mietelski, J.; Baeza, A.; Guillen, J.; Buzinny, M.; Tsigankov, N.; Gaca, P.; Jasińska, M.; Tomankiewicz, E. (2002). Plutonium and other alpha emitters in mushrooms from Poland, Spain and Ukraine. *Applied Radiation and Isotopes*, Vol. 56, pp. 717-729, ISSN 0969-8043.
- [72] Nikolova, I.; Johanson, K.; Dahlberg, A. (1997). Radiocaesium in fruitbodies and mycorrhizae in ectomycorrhizal fungi. *Journal of Environmental Radioactivity*, Vol.37, pp. 115-125, ISSN 0265-931X.
- [73] Nyholm, N. & Tyler G. (2000). Rubidium content of plant, fungi and animals closely reflects potassium and acidity conditions of forest soils. *Forest Ecology and Management*, Vol.134, pp. 89-96, ISSN 0378-1127.
- [74] Olsen, R.; Jøner, E.; Bakken. L. (1990). Soil fungi and the fate of radiocaesium in the soil ecosystem – a discussion of possible mechanisms involved in the radiocaesium accumulation in fungi, and the role of fungi as a Cs-sink in the soil.

- In: G. Desmet, P. Nassimbeni, & M. Belli (Eds.), *Transfer of radionuclides in natural and semi-natural environment*, pp. 657-663, ISBN 9781851665396, Luxemburg: Elsevier Applied Science.
- [75] Parekh, N.; Poskitt, J.; Dodd, B.; Potter, E.; Sanchez, A. (2008). Soil microorganisms determine the sorption of radionuclides within organic soil systems. *Journal of Environmental Radioactivity*, Vol.99, pp. 841-852, ISSN 0265-931X.
- [76] Pérez, A.; Farías, S.; Strobl, A.; Pérez, L.; López, C.; Piñeiro, A.; Roses, O.; Fajardo, M. (2007). Levels of essential and toxic elements in *Porphyra columbina* and *Ulva* sp. from San Jorge Gulf, Patagonia Argentina, *Science of the Total Environment*, Vol. 376, No. 1, pp. 51-59, ISSN 0048-9697.
- [77] Porter B. Orr. (1975). Available from:
<http://scholar.lib.vt.edu/ejournals/JARS/v30n3/v30n3-orr.htm>.
- [78] Radulescu, C.; Stihl, C.; Cimpoia, V. G.; Popescu, I. V.; Busuioc, G.; Gheboianu, A. I. (2011). Evaluation of heavy metals content in edible mushrooms by microwave digestion and Flame Atomic Absorption Spectrometry. *Scientific Study & Research, Chemistry & Chemical Engineering, Biotechnology, Food Industry*, Vol.12, No. 2, pp. 155-164, ISSN 1582540X.
- [79] Rafferty, B.; Brennan, M.; Dawson, D.; Dowding, D. (2000). Mechanisms of ^{137}Cs migration in coniferous forest soils. *Journal of Environmental Radioactivity*, Vol.48, pp. 131-143, ISSN 0265-931X.
- [80] Rafferty, B.; Dawson, D. & Kliashtorin, A. (1997). Decomposition in two pine forests: the mobilisation of ^{137}Cs and K from forest litter. *Soil Biology and Biochemistry*, Vol. 29, No. 11/12, pp. 1673-1681, ISSN 0038-0717.
- [81] Rappoldt, C.; Pieters, G.; Adema, E.; Baaijens, G.; Grootjans, A.; van Duijn, C. (2003). Buoyancy-driven flow in peat bogs as a mechanism for nutrient recycling. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 100, pp. 14937-14942, ISSN 1091-6490.
- [82] Read, D. & Perez-Moreno, J. (2003). Mycorrhizas and nutrient cycling in ecosystems – a journey towards relevance? *New Phytologist*, Vol. 157, pp. 475-492, ISSN 0028-646X.

- [83] Rodríguez-Navarro, A. (2000). Potassium transport in fungi and plants. *BBA – Biochimica et Biophysica Acta*, Vol. 1469, pp. 1-30, ISSN 1388-1981.
- [84] Rodushkin, I.; Engström, E.; Sörlin, D.; Baxter, D. (2008). Levels of inorganic constituents in raw nuts and seeds on the Swedish market. *Science of the Total Environment*, Vol. 39, No.2-3, pp. 290-304, ISSN 0048-9697.
- [85] Römmelt, R.; Hiersche, L.; Schaller, G.; Wirth, E. (1990). Influence of soil fungi (basidiomycetes) on the migration of ^{134}Cs + ^{137}Cs and ^{90}Sr in coniferous forest soils. In: G. Desmet, P. Nassimbeni and M. Belli, Editors, *Transfer of radionuclides in natural and seminatural environments*, pp. 152–160, ISBN 9781851665396, Elsevier, London.
- [86] Rosén, K.; Vinichuk, M.; Johanson, K. (2009). ^{137}Cs in a raised bog in central Sweden. *Journal of Environmental Radioactivity*, Vol. 100, pp. 534-539, ISSN 0265-931X.
- [87] Rosén, K.; Vinichuk, M.; Nikolova, I.; Johanson, K. (2011). Long-term effects of single potassium fertilization on ^{137}Cs levels in plants and fungi in a boreal forest ecosystem. *Journal of Environmental Radioactivity*, Vol. 102, pp. 178-184, ISSN 0265-931X.
- [88] Rühm, W.; Kammerer, L.; Hiersche, L.; Wirth, E. (1997). The $^{137}\text{Cs}/^{134}\text{Cs}$ ratio in fungi as an indicator of the major mycelium location in forest soil. *Journal of Environmental Radioactivity*, Vol.35, pp. 129-148, ISSN 0265-931X.
- [89] Rudawska, M. & Leski, N. (2005). Trace elements in fruiting bodies of ectomycorrhizal fungi growing in Scots pine (*Pinus sylvestris* L.) stands in Poland. *Science of the Total Environment*, Vol. 339, No.1-3, pp. 103-115, ISSN 0048-9697.
- [90] Rufyikiri, G.; Thiry, Y.; Wang, L.; Delvaux, B.; Declerck, S. (2002). Uranium uptake and translocation by the arbuscular mycorrhizal fungus, *Glomus intraradices*, under root-organ culture conditions. *New Phytologist*, Vol. 156, No. 2, pp. 275-28, ISSN 0028-646X.
- [91] Russell, E. (1988). *Soil conditions and plants growth*, Eleventh Edition, Longmans, ISBN 0582446775, London, England.

- [92] Rydin, H. & Clymo, R. (1989). Transport of carbon and phosphorus about Sphagnum. *Proceedings of Royal Society, Biol. Sci.*, Vol.237, pp. 63-84, ISSN 1471-2954, London.
- [93] Rydin, H. & Jeglum, J. (2006). *The biology of peatlands*, ISBN 978-0-19-852871-5, Oxford University Press.
- [94] Schmitt, J. A. & Meisch, H. U. (1985). Cadmium in mushrooms – distribution, growth effects and binding. *Trace Elements in Medicine*, Vol. 2, No.4, pp. 163-166, ISSN 0174-7371.
- [95] Seeger, R. & Schweinshaut, P. (1981). Vorkommen von caesium in höheren pilzen. *Science of the Total Environment*, Vol. 19, pp. 253-276, ISSN 0048-9697.
- [96] Skuterud, L.; Travnicova, I.; Balonov, M.; Strand, P.; Howard, B. (1997). Contribution of fungi to radiocaesium intake by rural populations in Russia. *Science of the Total Environment*, Vol. 193, No.3, pp. 237-242, ISSN 0048-9697.
- [97] Slekovec, M. & Irgolic, K. (1996). Uptake of arsenic my mushrooms from soil. *Chemical Speciation and Bioavailability*, Vol. 8, No. 3-4, pp. 67-73, ISSN 0954-2299.
- [98] Smith, S. & Read, D. (1997). *Mycorrhizal symbiosis* (2nd ed.), ISBN 0-12-652840-3, London: Academic Press.
- [99] Steiner, M.; Linkov, I.; Yoshida, S. (2002). The role of fungi in the transfer and cycling of radionuclides in forest ecosystems. *Journal of Environmental Radioactivity*, Vol. 58, pp. 217-241, ISSN 0265-931X.
- [100] Sybyrkyna, A. (2012). Accumulation of chrome by pine mushrooms Semipalatinsk Pryyrushja Republic Kazakhstan. Modern problems of science and education, Vol. 2. url: www.science-education.ru/102-5449.
- [101] Terada, H.; Shibata, H.; Kato, F.; Sugiyama, H. (1998). Influence of alkali elements on the accumulation of radiocesium by mushrooms. *Journal of Radioanalytical and Nuclear Chemistry*, Vol.235, No.1-2, pp. 195-200, ISSN 0236-5731.
- [102] Thomet, U.;Vogel, E.; Kr ähenbühl, U. (1999). The uptake of cadmium and zinc by mycelia and their accumulation in mycelia and fruiting bodies of edible

- mushrooms. *European Food Research and Technology*, Vol. 209, No.5, pp. 317-324, ISSN 1438-2377.
- [103] Tsukada, H.; Takeda, A.; Hisamatsu, S.; Inaba, J. Inequilibrium between Fallout ^{137}Cs and Stable Cs in Cultivated Soils. Radionuclides in Soils and Sediments, and their Transfer to Biota, In: *Proceedings of 18th World Congress of Soil Science*, 15 July 2006 Available from <http://crops.confex.com/crops/wc2006/techprogram/P17531.HTM>.
- [104] Turtiainen, T., Brunfeldt, M., Rasilainen, T., Skipperud, L., Valle, L., Popic, J. M., Roos, P., Sundell-Bergman, S., Ros n, K., Weimer, R. (2013). Doses from natural radioactivity in wild mushrooms and berries to the Nordic population. Interim Report from the NKS-B BERMUDA activity.
- [105] Nordisk Kernesikkerhedsforskning, 16 p. (NKS; No. 273). ISBN 978-87-7893-370-6.
- [106] Tyler, G. (1982). Accumulation and exclusion of metals in *Collybia peronata* and *Amaniuta rubescens*. *Transactions of British Mycological Society*, Vol. 79, pp. 239-241, ISSN 0007-1536.
- [107] van der Linden, M.; Vickery, E.; Charman, D.; van Geel, B. (2008). Effects of human impact and climate change during the last 350 years recorded in a Swedish raised bog deposit. *Palaeogeography, Palaeoclimatology, Palaeoecology*, Vol. 262, pp. 1-31, ISSN 0031-0182.
- [108] Vetter, J. (1994). Data on arsenic and cadmium contents of some common mushrooms. *Toxicon*, Vol. 82, No. 1, pp. 11-15, ISSN 0041-0101.
- [109] Vetter, J. (2005). Mineral composition of basidiomes of *Amanita* species, *Mycological Research*, Vol. 109, No.6, pp. 746-750, ISSN 0953-7562.
- [110] Villaverde, P.; Gondar, D.; Antelo, J.; Lpez, R.; Fiol, S.; Arce, F. (2009). Influence of pH on copper, lead and cadmium binding by an ombrotrophic peat. *European Journal of Soil Science*, Vol. 60, No. 3, pp. 377-385, ISSN 1351-0754.
- [111] Vinichuk, M. & Johanson, K. (2003). Accumulation of ^{137}Cs by fungal mycelium in forest ecosystems of Ukraine. *Journal of Environmental Radioactivity*, Vol. 64, pp. 27-43, ISSN 0265-931X.

- [112] Vinichuk, M.; Johanson, K.; Taylor, A. (2004). ^{137}Cs in the fungal compartment of Swedish forest soils. *Science of the Total Environment*, Vol. 323, pp. 243-251, ISSN 0048-9697.
- [113] Vinichuk, M.; Johanson, K.; Rosén, K.; Nilsson, I. (2005). Role of fungal mycelium in the retention of radiocaesium in forest soils. *Journal of Environmental Radioactivity*, Vol. 78, pp. 77-92, ISSN 0265-931X.
- [114] Vinichuk, M.; Johanson, K.; Rydin, H.; Rosén, K. (2010a). The distribution of ^{137}Cs , K, Rb and Cs in plants in a Sphagnum-dominated peatland in eastern central Sweden. *Journal of Environmental Radioactivity*, Vol. 101, pp. 170-176, ISSN 0265-931X.
- [115] Vinichuk, M.; Taylor, A.; Rosén, K.; Johanson, K. (2010b). Accumulation of potassium, rubidium and cesium (^{133}Cs and ^{137}Cs) in various fractions of soil and fungi in a Swedish forest. *Science of the Total Environment*, Vol. 408, pp. 2543-2548, ISSN 0048-9697
- [116] Vinichuk, M.; Rosén, K.; Johanson, K.; Dahlberg, A. (2011a). Correlations between potassium, rubidium and cesium (^{133}Cs and ^{137}Cs) in sporocarps of *Suillus variegatus* in a Swedish boreal forest. *Journal of Environmental Radioactivity*, Vol. 102, No.4, pp. 386-392, ISSN 0265-931X.
- [117] Vinichuk, M.; Dahlberg, A.; Rosén, K. (2011b). Cesium (^{137}Cs and ^{133}Cs), Potassium and Rubidium in Macromycete Fungi and Sphagnum Plants, Radioisotopes – Applications in Physical Sciences, Nirmal Singh (Ed.), ISBN 978-953-307-510-5, InTech. (In English), pp. 279-310.
- [118] Vinichuk, M. (2012a). Chromium and nickel in soil fractions and certain macromycete species of boreal forest ecosystems. *Visnyk of Zaporizhzhya National University. Biological Sciences*, Vol.3, pp. 103-110. (ukr).
- [119] Vinichuk, M. (2012b). Selected Metals in Various Fractions of Soil and Fungi in a Swedish Forest. *ISRN Ecology*, Vol. 2012, ID 521582, 7 pages, ISSN 2090-4622.
- [120] Vinichuk, M. (2012c). Thorium and uranium in soil fractions and certain macromycete species in boreal forest ecosystems. *Visnyk of Dnipropetrovsk University. Biology. Ecology*, Vol. 20(2), pp. 10-17. (ukr).

- [121] Vinichuk, M. (2013). Copper, zinc, and cadmium in various fractions of soil and fungi in a Swedish forest. *Journal of Environmental Science and Health, Part A*, Vol.48, No.8, pp. 980-987.
- [122] Vinichuk, M.; Ros n, K.; Dahlberg, A. (2013). ¹³⁷Cs in fungal sporocarps in relation to vegetation in a bog, pine swamp and forest along a transect. *Chemosphere*, Vol. 90, No.2, pp. 713-720, ISSN 0045-6535.
- [123] Wallace, A. (1970). Monovalent-ion carrier effects on transport of ⁸⁶Rb and ¹³⁷Cs into bush bean plants. *Plant and Soil*, Vol.32, pp. 526-520, ISSN 0032-079X.
- [124] White, P. & Broadley, M. (2000). Mechanisms of caesium uptake by plants. *New Phytologist*, Vol. 147, pp. 241-256, ISSN 0028-646X.
- [125] Yoshida, S. & Muramatsu, Y. (1998). Concentration of alkali and alkaline earth elements in mushrooms and plants collected in a Japanese pine forest, and their relationship with ¹³⁷Cs. *Journal of Environmental Radioactivity*, Vol.41, No.2, pp. 183-205, ISSN 0265-931X.
- [126] Zichner, A. (2000). Cesium uptake by fungi in the presence of illite. Internet resource: <http://dx.doi.org/10.3929/ethz-a-003906656>.
- [127] Zurera-Cosano, G.; Rincon-Leon, F.; Pozo-Lora, R. (1988). Lead and Cadmium content of some edible mushrooms. *Journal of Food Quality*, Vol. 10, No.5, pp. 311-317, ISSN 0146-9428.



Mykhailo Vinichuk is Professor in Ecology, Zhytomyr State Technological University, Zhytomyr, Ukraine. His research interests are behavior of radionuclides in both natural (forest, peatland) and agricultural ecosystems with special emphasis on the role of ectomycorrhizal fungi in radionuclides transfer.



Klas Rosén is Associate Professor at Department of Soil & Environment, Swedish University of Agricultural Sciences SLU, Sweden. His research interests are both in terrestrial radioecology and radiation safety with various countermeasures in agricultural, forest and urban areas.

To order additional copies of this book, please contact:
Science Publishing Group
book@sciencepublishinggroup.com
www.sciencepublishinggroup.com

